

THE CYCLOTOMIC BMW ALGEBRA ASSOCIATED WITH THE TWO STRING TYPE B BRAID GROUP

STEWART WILCOX AND SHONA YU

ABSTRACT. The cyclotomic Birman-Murakami-Wenzl (or BMW) algebras $\mathcal{B}_n^k$, introduced by R. Häring-Oldenburg, are extensions of the cyclotomic Hecke algebras of Ariki-Koike, in the same way as the BMW algebras are extensions of the Hecke algebras of type A . In this paper we focus on the case $n = 2$, producing a basis of $\mathcal{B}_2^k$ and constructing its left regular representation.

1. INTRODUCTION

Motivated by knot theory associated to the Artin braid group of type A and the Kauffman link polynomial, Murakami [5] and Birman and Wenzl [2] defined what are now known as the BMW algebras. The Artin braid group relations of type A appear in the defining relations of the BMW algebras. Also, the Hecke algebra of the symmetric group (the Coxeter group of type A) appears naturally as a quotient of this algebra. Motivated by type B knot theory and the Ariki-Koike algebras [1] (also known as the cyclotomic Hecke algebras of type $G(k, 1, n)$), Häring-Oldenburg [4] introduced the “cyclotomic BMW algebras” $\mathcal{B}_n^k$, of which the Ariki-Koike algebras appear naturally as a quotient in a similar fashion. In this paper we study in detail the algebra $\mathcal{B}_2^k$, with the hope of utilising our results to study the algebra for higher n .

A natural question to address is whether $\mathcal{B}_2^k$ is always free as a module over our ground ring R . Our main result is that if our parameters $A_0, \dots, A_{k-1}, q_0, \dots, q_{k-1}, q, \lambda$ are admissible (see Definition 2) then the algebra $\mathcal{B}_2^k$ is R -free of rank $3k^2$.

In Section 2, we establish these “admissibility” conditions (see Definition 2) and construct a $\mathcal{B}_2^k$ -module V of rank k (see Lemma 3). Then in Section 3, using this rank k module, we are able to construct the regular representation of $\mathcal{B}_2^k$ and provide an explicit basis of the algebra. These results are stated but incompletely proved in Häring-Oldenburg [4]; specifically, additional arguments are needed to prove Lemma 25. In the present paper we take a slightly different approach and the arguments we offer correct this problem.

The authors thank Bob Howlett for his helpful suggestions during the preparation of this manuscript.

2. THE CYCLOTOMIC BMW ALGEBRAS

Definition 1. Fix natural numbers n and k . Let R be a commutative ring containing units $A_0, p_0, \dots, p_{k-1}, q, \lambda$ and further elements $A_1, \dots, A_{k-1}$ such that $\lambda - \lambda^{-1} = \delta(1 - A_0)$ holds, where $\delta = q - q^{-1}$. The **cyclotomic BMW algebra** $\mathcal{B}_n^k(R)$ is defined to be the unital associative R -algebra generated by $Y^{\pm 1}, X_1^{\pm 1}, \dots, X_{n-1}^{\pm 1}$ and $e_1, \dots, e_{n-1}$ subject to the following relations:

$$\begin{aligned}
X_i - X_i^{-1} &= \delta(1 - e_i) \\
X_i X_j &= X_j X_i \quad \text{for } |i - j| \geq 2 \\
X_i X_{i+1} X_i &= X_{i+1} X_i X_{i+1} \\
X_i e_j &= e_j X_i \quad \text{for } |i - j| \geq 2 \\
e_i e_j &= e_j e_i \quad \text{for } |i - j| \geq 2 \\
X_i e_i &= e_i X_i = \lambda e_i \\
e_i X_{i+1} e_i &= \lambda^{-1} e_i \\
X_i X_{i+1} e_i &= e_{i+1} e_i = e_{i+1} X_i X_{i+1} \\
e_i e_{i+1} e_i &= e_i \\
\prod_{i=0}^{k-1} (Y - p_i) &= 0 \\
X_1 Y X_1 Y &= Y X_1 Y X_1 \\
Y X_i &= X_i Y \quad \text{for } i > 1 \\
Y X_1 Y e_1 &= \lambda^{-1} e_1 = e_1 Y X_1 Y \\
e_1 Y^m e_1 &= A_m e_1 \quad \text{for } 0 \leq m \leq k-1.
\end{aligned}$$

If the k^{th} order relation $\prod_{i=0}^{k-1} (Y - p_i) = 0$ is omitted one obtains the affine BMW algebra, studied by Goodman and Hauschild in [3]. Thus the cyclotomic BMW algebra is a quotient of the affine BMW algebra.

In this paper we shall focus our attention on the algebra $\mathcal{B}_2^k(R)$. We simplify our notation by omitting the index 1 of X_1 and e_1 . Specifically, $\mathcal{B}_2^k(R)$ is the unital associative R -algebra generated by $Y^{\pm 1}, X^{\pm 1}$ and e subject to the following relations:

$$\prod_{i=0}^{k-1} (Y - p_i) = 0 \tag{1}$$

$$X - X^{-1} = \delta(1 - e) \tag{2}$$

$$XYXY = YXYX \tag{3}$$

$$Xe = \lambda e = eX \tag{4}$$

$$YXYe = \lambda^{-1} e = eYXY \tag{5}$$

$$eY^m e = A_m e \quad \text{for } 0 \leq m \leq k-1. \tag{6}$$

Let $q_0, \dots, q_{k-1}$ be the signed elementary symmetric polynomials in $p_0, \dots, p_{k-1}$ so that

$$Y^k = \sum_{i=0}^{k-1} q_i Y^i.$$

Observe that $q_0 = (-1)^{k-1} \prod_i p_i$ is invertible. Define $q_k = -1$. Then $\sum_{l=0}^k q_l Y^l = 0$ and the inverse of Y may then be expressed as a linear combination of nonnegative powers of Y as follows:

$$Y^{-1} = -q_0^{-1} \sum_{i=0}^{k-1} q_{i+1} Y^i.$$

Remark:

1) Our proofs apply unchanged if the relation $\prod_{i=0}^{k-1} (Y - p_i) = 0$ is replaced by $Y^k = \sum_{i=0}^{k-1} q_i Y^i$, without any assumption that the polynomial $\sum_{j=0}^k q_j x^j$ factorizes in R . All that we require is that q_0 be invertible.

2) There is an anti-involution $*$ of $\mathcal{B}_2^k(R)$ such that

$$Y^* = Y, \quad X^* = X \quad \text{and} \quad e^* = e. \quad (7)$$

Let $\mathfrak{h}_2^k(R)$ denote the unital associative R -algebra with generators $T_0^{\pm 1}, T_1^{\pm 1}$ and relations

$$\begin{aligned} T_0 T_1 T_0 T_1 &= T_1 T_0 T_1 T_0 \\ \prod_{i=0}^{k-1} (T_0 - p_i) &= 0 \\ T_1^2 &= \delta T_1 + 1. \end{aligned}$$

Here $\mathfrak{h}_2^k(R)$ is essentially the Ariki-Koike algebra [1], with the parameters modified. It is clearly a quotient of $\mathcal{B}_2^k(R)$, and indeed $\mathcal{B}_2^k/I \cong \mathfrak{h}_2^k$ as R -algebras, where I is the two-sided ideal generated by e in $\mathcal{B}_2^k(R)$.

Using the relations of the algebra and the relation

$$\begin{aligned} XY^l e &= Y^{-1} X^{-1} X Y X Y^l e \\ &\stackrel{(3)}{=} Y^{-1} X^{-1} Y^l X Y X e \\ &\stackrel{(4),(5)}{=} Y^{-1} X^{-1} Y^{l-1} e \\ &\stackrel{(2),(6)}{=} Y^{-1} X Y^{l-1} e - \delta Y^{l-2} e + \delta A_{l-1} Y^{-1} e, \end{aligned}$$

it is straightforward to show the left ideal of $\mathcal{B}_2^k$ generated by e is the span of $\{Y^i e \mid 0 \leq i \leq k-1\}$.

As a consequence of the results in Goodman and Hauschild [3], the set $\{Y^i e \mid i \in \mathbb{Z}\}$ is linearly independent in the affine BMW algebra, and so it seems natural to expect that the set $\{Y^i e \mid 0 \leq i \leq k-1\}$ will be linearly independent in the cyclotomic BMW algebra. However, for this to hold we must impose additional restrictions on our

parameters $A_0, \dots, A_{k-1}, q_0, \dots, q_{k-1}, q, \lambda$. We say that our choice of parameters $A_0, \dots, A_{k-1}, q_0, \dots, q_{k-1}, q, \lambda$ is *admissible* if these conditions are satisfied and, additionally, $q - q^{-1}$ is not a zero divisor. The precise definition is given below.

3. CONSTRUCTION OF V

Under these admissibility conditions, we construct a free $\mathcal{B}_2^k(R)$ -module V of rank k isomorphic to the R -span of $\{Y^i e \mid 0 \leq i \leq k-1\}$, hence proving the $\{Y^i e \mid 0 \leq i \leq k-1\}$ are indeed linearly independent.

Let V be the free R -module of rank k with basis $v_0, v_1, \dots, v_{k-1}$. Define a linear map $\bar{Y} : V \rightarrow V$ by

$$\bar{Y}v_i = v_{i+1}, \quad \text{for } 0 \leq i \leq k-2, \quad (8)$$

$$\bar{Y}v_{k-1} = \sum_{i=0}^{k-1} q_i v_i. \quad (9)$$

Since q_0 is invertible, this guarantees that $\bar{Y}$ is invertible, with inverse

$$\bar{Y}^{-1}v_i = v_{i-1}, \quad \text{for } 1 \leq i \leq k-1, \quad (10)$$

$$\bar{Y}^{-1}v_0 = -q_0^{-1} \sum_{i=0}^{k-1} q_{i+1} v_i. \quad (11)$$

Also $\bar{Y}^i v_0 = v_i$ for $i = 0, \dots, k-1$. Define $v_s = \bar{Y}^s v_0$ for all integers s . The definition of $\bar{Y}v_{k-1}$ gives

$$\sum_{l=0}^k q_l \bar{Y}^l v_0 = 0.$$

For any integer i , applying $\bar{Y}^i$ to this gives

$$\sum_{l=0}^k q_l \bar{Y}^{l+i} v_0 = 0. \quad (12)$$

Also $\bar{Y}^l$ is invertible for any integer l and, as $\{v_0, v_1, \dots, v_{k-1}\}$ is a basis for V , the set

$$\{\bar{Y}^l v_0, \bar{Y}^l v_1, \dots, \bar{Y}^l v_{k-1}\} = \{v_l, v_{l+1}, \dots, v_{l+k-1}\} \quad (13)$$

is also a basis for V .

Now let us define linear maps $\bar{X}, \bar{E} : V \rightarrow V$ by

$$\bar{X}v_0 = \lambda v_0, \quad (14)$$

$$\bar{X}v_1 = \lambda^{-1} \bar{Y}^{-1} v_0, \quad (15)$$

$$\bar{X}v_i = \bar{Y}^{-1} \bar{X}v_{i-1} - \delta v_{i-2} + \delta A_{i-1} \bar{Y}^{-1} v_0, \quad \text{for } 2 \leq i \leq k-1, \quad (16)$$

$$\bar{E}v_i = A_i v_0, \quad \text{for } 0 \leq i \leq k-1. \quad (17)$$

Since $\lambda^{-1} = \lambda - \delta + \delta A_0$ in R , substituting $i = 1$ into (16) reproduces (15). Also, note that the image of the map $\bar{E}$ is $\text{Im}(\bar{E}) = \langle v_0 \rangle$, where $\langle S \rangle$ denotes the R -submodule spanned by the set S . Furthermore, let us denote by $\bar{W}$ the map $\bar{X} - \delta + \delta \bar{E}$. Thus (15) and (16) become

$$\bar{X}v_i = \bar{Y}^{-1}\bar{W}v_{i-1}, \quad \text{for } 1 \leq i \leq k-1. \quad (18)$$

Our aim is to show that V is actually a $\mathcal{B}_2^k$ -module, where the action of the generators Y , X and e are given by the maps $\bar{Y}$, $\bar{X}$ and $\bar{E}$, respectively. In order to prove this, we require $\bar{Y}^{-1}\bar{W}\bar{Y}^{-1} = \bar{X}$. By equations (10) and (18), this relation automatically holds on $v_1, \dots, v_{k-1}$, so we need only ensure that

$$(\bar{Y}^{-1}\bar{W}\bar{Y}^{-1} - \bar{X})v_0 = 0.$$

For convenience, we write the left hand side relative to the basis $\{v_0, v_{-1}, \dots, v_{1-k}\}$. Let

$$(\bar{Y}^{-1}\bar{W}\bar{Y}^{-1} - \bar{X})v_0 = \alpha v_0 + \sum_{l=1}^{k-1} h_l v_{-l},$$

where α and h_l are elements of R .

Definition 2. Let R be as in Definition 1. The family of parameters $(A_0, \dots, A_{k-1}, q_0, \dots, q_{k-1}, q, \lambda)$ is called **admissible** if the following three properties are satisfied:

- (a) $\delta = q - q^{-1}$ is not a zero divisor;
- (b) $\lambda - \lambda^{-1} = \delta(1 - A_0)$;
- (c) $h_1 = \dots = h_{k-1} = \alpha = 0$, where the h_l and α are as defined above.

It can be shown that the admissibility conditions are given by the following:

$$\alpha = \begin{cases} q_0^{-2}\lambda^{-1} - \lambda & \text{if } k \text{ is odd,} \\ q_0^{-2}\lambda^{-1} - \lambda - \delta q_0^{-1} & \text{if } k \text{ is even,} \end{cases}$$

and

$$h_l = -\lambda^{-1}(q_0^{-1}q_l + q_0^{-2}q_{k-l}) - \delta q_0^{-1} \left[\sum_{r=1}^{k-l} q_{r+l} A_r - \sum_{i=\max(l+1, m)}^{\lfloor \frac{l+k}{2} \rfloor} q_{2i-l} + \sum_{i=\lceil \frac{l}{2} \rceil}^{\min(l, m-1)} q_{2i-l} \right].$$

In the case when δ is invertible, these conditions may be viewed as a triangular system of equations for $A_0, \dots, A_{k-1}$. It is important to remark here that our definition of admissibility differs from the definitions of “potentially admissible” and “admissible” in Häring-Oldenburg [4]. However, the above equations coincide with those obtained by Häring-Oldenburg. Also, note that R need not be an integral domain.

Henceforth assume that R is as in Definition 1 and the parameters $A_0, \dots, A_{k-1}, q_0, \dots, q_{k-1}, q, \lambda$ are admissible. Denote $\mathcal{B}_2^k(R)$ simply by $\mathcal{B}_2^k$.

Then, by our argument above, $\bar{Y}^{-1}\bar{W}\bar{Y}^{-1} = \bar{X}$ holds on all of V and hence rearranging gives

$$\bar{Y}\bar{X}\bar{Y} = \bar{W} = \bar{X} - \delta + \delta\bar{E}.$$

Thus $\bar{Y}\bar{X}\bar{Y}\bar{X} - \bar{X}\bar{Y}\bar{X}\bar{Y} = [\bar{Y}\bar{X}\bar{Y}, \bar{X}] = [\delta\bar{E}, \bar{X}]$, where $[\cdot, \cdot]$ denotes the standard commutator of two maps.

Since $\text{Im}(\bar{E}) = \langle v_0 \rangle$ and $\bar{X}v_0 = \lambda v_0$ by definition,

$$\text{Im}([\delta\bar{E}, \bar{X}]) \subseteq \langle v_0 \rangle. \quad (19)$$

Let $N = \bar{Y}\bar{X}\bar{Y}\bar{X} - 1$. Then

$$\begin{aligned} [N, \bar{Y}] &= N\bar{Y} - \bar{Y}N \\ &= \bar{Y}\bar{X}\bar{Y}\bar{X}\bar{Y} - \bar{Y} - \bar{Y}\bar{Y}\bar{X}\bar{Y}\bar{X} + \bar{Y} \\ &= \bar{Y}(\bar{X}\bar{Y}\bar{X}\bar{Y} - \bar{Y}\bar{X}\bar{Y}\bar{X}) \\ &= -\bar{Y}[\delta\bar{E}, \bar{X}] \end{aligned}$$

and

$$[N, \bar{Y}^{-1}] = -\bar{Y}^{-1}[N, \bar{Y}]\bar{Y}^{-1} = [\delta\bar{E}, \bar{X}]\bar{Y}^{-1}.$$

Therefore, by (19),

$$\text{Im}([N, \bar{Y}]) \subseteq \langle v_1 \rangle \quad \text{and} \quad \text{Im}([N, \bar{Y}^{-1}]) \subseteq \langle v_0 \rangle. \quad (20)$$

Observe that

$$\bar{Y}\bar{X}\bar{Y}\bar{X}v_0 = \lambda\bar{Y}\bar{X}\bar{Y}v_0 = \lambda\bar{Y}\bar{X}v_1 = \lambda\bar{Y}(\lambda^{-1}\bar{Y}^{-1}v_0) = v_0$$

and

$$\bar{Y}\bar{X}\bar{Y}\bar{X}v_1 = \bar{Y}\bar{X}\bar{Y}(\lambda^{-1}\bar{Y}^{-1}v_0) = \lambda^{-1}\bar{Y}\bar{X}v_0 = \lambda^{-1}\bar{Y}(\lambda v_0) = v_1,$$

hence $Nv_0 = Nv_1 = 0$.

Lemma 1.

$$Nv_l \in \langle v_1, v_2, \dots, v_{l-1} \rangle, \quad \text{for all } l \geq 1, \quad (21)$$

and

$$Nv_{-m} \in \langle v_0, v_{-1}, \dots, v_{-(m-1)} \rangle, \quad \text{for all } m \geq 0. \quad (22)$$

Proof. We have already established $Nv_0 = Nv_1 = 0$ above. To prove the first assertion, we argue by induction on l . Assume that $l \geq 2$ and $Nv_{l-1} \in \langle v_1, v_2, \dots, v_{l-2} \rangle$. Then

$$Nv_l = [N, \bar{Y}]v_{l-1} + \bar{Y}Nv_{l-1} \in \langle v_1 \rangle + \langle v_2, \dots, v_{l-1} \rangle,$$

by (20) and the inductive hypothesis. Thus $Nv_l \in \langle v_1, \dots, v_{l-1} \rangle$ for all $l \geq 1$.

Similarly, we prove the second assertion by induction on m . Assume $m \geq 1$ and $Nv_{-(m-1)} \in \langle v_0, v_{-1}, \dots, v_{-(m-2)} \rangle$. Then

$$\begin{aligned} Nv_{-m} &= [N, \bar{Y}^{-1}]v_{-(m-1)} + \bar{Y}^{-1}Nv_{-(m-1)} \\ &\in \langle v_0 \rangle + \langle v_{-1}, v_{-2}, \dots, v_{-(m-1)} \rangle, \end{aligned}$$

by (20) and the inductive hypothesis. So $Nv_{-m} \in \langle v_0, v_{-1}, \dots, v_{-(m-1)} \rangle$ for all $m \geq 0$. $\square$

Lemma 2. $\bar{Y}\bar{X}\bar{Y}\bar{X}$ is the identity map on V .

Proof. We are required to show that $Nv_i = 0$ for $i = 0, 1, \dots, k-1$. We proceed by induction on i . The cases $i = 0$ and $i = 1$ have been established above. Suppose that $2 \leq i \leq k-1$ and

$$Nv_0 = Nv_1 = \dots = Nv_{i-1} = 0. \quad (23)$$

Since $\{v_{i-1}, v_{i-2}, \dots, v_{i-k}\}$ is a basis for V , by (13), we have that

$$v_i \in \langle v_{i-1}, v_{i-2}, \dots, v_0, v_{-1}, \dots, v_{i-k} \rangle.$$

Together with our inductive hypothesis (23), this implies

$$\begin{aligned} Nv_i &\in N\langle v_0, v_{-1}, \dots, v_{i-k} \rangle \\ &\subseteq \langle v_0, v_{-1}, \dots, v_{i-k+1} \rangle, \text{ by (22).} \end{aligned}$$

However (21) states that $Nv_i \in \langle v_1, v_2, \dots, v_{i-1} \rangle$. Again using that $\{v_{i-1}, v_{i-2}, \dots, v_{i-k+1}, v_{i-k}\}$ forms a basis for V ,

$$\langle v_1, v_2, \dots, v_{i-1} \rangle \cap \langle v_0, v_{-1}, \dots, v_{i-k+1} \rangle = 0.$$

Therefore $Nv_i = 0$. So by induction on i , we have proved that $Nv_i = 0$ for all $0 \leq i \leq k-1$, as required. $\square$

Lemma 3. (cf. Lemma 25 of Haring-Oldenburg [4])

Suppose the choice of parameters in R satisfy the admissibility conditions. Then the following relations hold on V :

$$\sum_{l=0}^k q_l \bar{Y}^l = 0 \quad (24)$$

$$\bar{X}\bar{W} = \bar{W}\bar{X} = 1 \quad (25)$$

$$\bar{X}\bar{E} = \lambda \bar{E} = \bar{E}\bar{X} \quad (26)$$

$$\bar{Y}\bar{X}\bar{Y}\bar{X} = \bar{X}\bar{Y}\bar{X}\bar{Y} \quad (27)$$

$$\bar{E}\bar{Y}^m \bar{E} = A_m \bar{E}, \text{ for all } 0 \leq m \leq k-1, \quad (28)$$

$$\bar{E}\bar{Y}\bar{X}\bar{Y} = \bar{Y}\bar{X}\bar{Y}\bar{E} = \lambda^{-1} \bar{E} \quad (29)$$

Furthermore V is a $\mathcal{B}_2^k(R)$ -module, where the actions of Y , X , X^{-1} and e on V are given by the maps $\bar{Y}$, $\bar{X}$, $\bar{W}$ and $\bar{E}$, respectively.

Proof. The k^{th} order relation (24) is immediate from (12). As a consequence of Lemma 2,

$$\overline{W}\overline{X} = \overline{Y}\overline{X}\overline{Y}\overline{X} = 1.$$

Clearly, this implies

$$\overline{X}\overline{W} = \overline{X}\overline{Y}\overline{X}\overline{Y} = Y^{-1}(\overline{Y}\overline{X}\overline{Y}\overline{X})\overline{Y} = Y^{-1}(1)\overline{Y} = 1.$$

Hence we have proved (25) and (27). Moreover,

$$[\delta\overline{E}, \overline{X}] = [\overline{W}, \overline{X}] = 0.$$

Since δ is not a zero divisor in R , by the definition of admissibility, and since $\text{End}_R(V)$ is a free R -module, it follows that $\overline{X}$ commutes with $\overline{E}$. Thus using (17) and (14),

$$\overline{E}\overline{X} = \overline{X}\overline{E} = \lambda\overline{E},$$

proving (26). Equation (28) follows easily from (8) and (17). Furthermore, since $\overline{Y}\overline{X}\overline{Y} = \overline{W}$, (25) and (26) imply (29). The last assertion of Lemma 3 now follows immediately. $\square$

Theorem 3.1. *The map $\varphi : V \rightarrow \text{span}_R\{Y^i e \mid 0 \leq i \leq k-1\}$ which maps v_i to $Y^i e$ defines a $\mathcal{B}_2^k(R)$ -module isomorphism.*

Proof. Let $U = \text{span}_R\{Y^i e \mid 0 \leq i \leq k-1\}$. It is clear that $\varphi : V \rightarrow U$ is a surjective R -module homomorphism. Recall that U is a left ideal in $\mathcal{B}_2^k$. We can therefore define a $\mathcal{B}_2^k$ -module homomorphism $\psi : U \rightarrow V$ by

$$\psi(a) = a(A_0^{-1}v_0), \quad \text{for all } a \in U.$$

Then

$$\psi(\varphi(v_i)) = Y^i e(A_0^{-1}v_0) = Y^i v_0 = v_i$$

for $0 \leq i \leq k-1$, so that $\psi\varphi$ is the identity. Since φ is surjective, it follows that ψ and φ are inverses. Therefore they are both $\mathcal{B}_2^k$ -module isomorphisms. $\square$

Hence, as a direct consequence of Theorem 3.1, the set $\{Y^i e \mid 0 \leq i \leq k-1\}$ is linearly independent. A more direct proof of this is given as follows. Suppose $\sum_{i=0}^{k-1} \alpha_i Y^i e = 0$, where $\alpha_i \in R$. Considering the action of both sides on v_0 gives

$$A_0 \sum_{i=0}^{k-1} \alpha_i v_i = 0.$$

But A_0 is invertible and $v_0, \dots, v_{k-1}$ are linearly independent over R , so each α_i must be 0.

4. CONSTRUCTING THE REGULAR REPRESENTATION OF $\mathcal{B}_2^k$

In this section, under the assumptions of admissibility, we shall prove that $\mathcal{B}_2^k(R)$ is a free R -module of rank $3k^2$ and provide an explicit basis of $\mathcal{B}_2^k(R)$; consequently, we show that $\{Y^i e Y^j \mid i, j = 0, \dots, k-1\}$ is a basis of the two-sided ideal of $\mathcal{B}_2^k(R)$ generated by e .

Recall that $\{v_i \mid r \leq i < r+k\}$ forms a basis for V for any integer r . Therefore, the R -module $\Xi_0 = V \otimes_R V$ has basis

$$\{v_{ij} \mid r \leq i < r+k \text{ and } s \leq j < s+k\}$$

for any integers r and s , where $v_{ij} = v_i \otimes v_j$. In particular, Ξ_0 is R -free of rank k^2 . The anti-involution $*$ permits us to make $\Xi_0 = V \otimes V$ into a $(\mathcal{B}_2^k, \mathcal{B}_2^k)$ -bimodule with left action satisfying

$$a(v \otimes v') = av \otimes v'$$

and right action satisfying

$$(v \otimes v')b = v \otimes (b^* v'),$$

for all $a, b \in \mathcal{B}_2^k$ and $v, v' \in V$. In particular, for each integer j there is a $\mathcal{B}_2^k$ -module isomorphism from V to $\langle v_{ij} \mid i \in \mathbb{Z} \rangle$ taking v_i to v_{ij} for all i .

Let Ξ_1 and Ξ_2 be R -modules isomorphic to Ξ_0 , with isomorphisms $\xi_1 : \Xi_0 \rightarrow \Xi_1$ and $\xi_2 : \Xi_0 \rightarrow \Xi_2$. Define

$$\Xi = \Xi_0 \oplus \Xi_1 \oplus \Xi_2,$$

so that Ξ is R -free of rank $3k^2$. For all integers i and j , define $u_{ij} = \xi_1(v_{ij}) \in \Xi_1$ and $w_{ij} = \xi_2(v_{ij}) \in \Xi_2$. Now (12) gives

$$\left(\sum_{l=0}^k q_l v_{i+l} \right) \otimes v_j = v_i \otimes \left(\sum_{l=0}^k q_l v_{j+l} \right) = 0.$$

Expanding and applying ξ_1 and ξ_2 , we obtain

$$\begin{aligned} \sum_{l=0}^k q_l v_{i+l,j} &= \sum_{l=0}^k q_l u_{i+l,j} = \sum_{l=0}^k q_l w_{i+l,j} = 0 \\ \sum_{l=0}^k q_l v_{i,j+l} &= \sum_{l=0}^k q_l u_{i,j+l} = \sum_{l=0}^k q_l w_{i,j+l} = 0, \end{aligned} \tag{30}$$

for any integers i and j .

In order to give Ξ a $\mathcal{B}_2^k$ -module structure, we shall define linear maps in $\text{End}_R(\Xi)$, again denoted $\bar{Y}$, $\bar{X}$ and $\bar{E}$, and show they satisfy the defining relations of $\mathcal{B}_2^k$. We already have a left $\mathcal{B}_2^k$ -module structure on Ξ_0 , so we define $\bar{Y}$, $\bar{X}$ and $\bar{E}$ to act on Ξ_0 as Y , X and e , respectively.

Thus it follows that

$$\bar{Y}v_{ij} = v_{i+1,j}, \quad (31)$$

and

$$\begin{aligned} \bar{X}v_{0j} &= \lambda v_{0j}, \\ \bar{X}v_{ij} &= Y^{-1}Xv_{i-1,j} - \delta v_{i-2,j} + \delta A_{i-1}Y^{-1}v_{0j}, \quad \text{for } 1 \leq i \leq k-1, \\ \bar{E}v_{ij} &= A_i v_{0j}, \quad \text{for } 0 \leq i \leq k-1. \end{aligned}$$

We want Ξ to be isomorphic to the regular representation with v_{ij} , u_{ij} and w_{ij} corresponding to $Y^i e Y^j$, $Y^i X Y^j$ and $X Y^i X Y^j$, respectively. Motivated by this, we make the following definitions for $i, j \in \{0, \dots, k-1\}$:

$$\bar{E}u_{ij} = v_{00}Y^iXY^j \quad (32)$$

$$\bar{E}w_{ij} = v_{00}XY^iXY^j = \lambda \bar{E}u_{ij} \quad (33)$$

$$\bar{X}u_{ij} = w_{ij} \quad (34)$$

$$\bar{X}w_{ij} = u_{ij} + \delta w_{ij} - \delta \lambda \bar{E}u_{ij} \quad (35)$$

$$\bar{Y}u_{ij} = u_{i+1,j} \quad (36)$$

$$\bar{Y}w_{ij} = \bar{W}u_{i,j+1} + \delta u_{1,i+j} - \delta YXv_{ij}, \quad (37)$$

where, as in the construction of the $\mathcal{B}_2^k$ -module V , $\bar{W}$ denotes the linear map $\bar{X} - \delta + \delta \bar{E}$. Note that (32) and (33) make use of the right $\mathcal{B}_2^k$ -module structure of Ξ_0 . Since v_{ij} , u_{ij} and w_{ij} satisfy the k -th order relations (30), an easy argument shows that (31)–(37) are satisfied for any integers i and j .

Lemma 4. *For any integer i ,*

$$\bar{Y}w_{1i} = w_{1,i+1}. \quad (38)$$

Proof. Because $YXYv_0 = X^{-1}v_0 = \lambda^{-1}v_0$, we have

$$YXYv_{0i} = \lambda^{-1}v_{0i} \quad \text{and} \quad v_{00}YXY = \lambda^{-1}v_{00}.$$

Thus

$$\begin{aligned} \bar{Y}w_{1i} &\stackrel{(37)}{=} \bar{W}u_{1,i+1} + \delta u_{1,i+1} - \delta YXv_{1i} \\ &\stackrel{(31)}{=} (\bar{X} - \delta + \delta \bar{E})u_{1,i+1} + \delta u_{1,i+1} - \delta YXYv_{0i} \\ &\stackrel{(32)}{=} \bar{X}u_{1,i+1} + \delta v_{00}YXY^{i+1} - \delta \lambda^{-1}v_{0i} \\ &\stackrel{(34)}{=} w_{1,i+1} + \delta \lambda^{-1}v_{00}Y^i - \delta \lambda^{-1}v_{00}Y^i \\ &= w_{1,i+1}, \end{aligned}$$

as required. $\square$

Theorem 4.1. *The maps $\bar{X}, \bar{W}, \bar{Y}, \bar{E} : \Xi \rightarrow \Xi$ defined above satisfy the following identities:*

$$\bar{X}\bar{W} = \bar{W}\bar{X} = 1 \quad (39)$$

$$\bar{X}\bar{E} = \bar{E}\bar{X} = \lambda\bar{E} \quad (40)$$

$$\bar{X}\bar{Y}\bar{X}\bar{Y} = \bar{Y}\bar{X}\bar{Y}\bar{X} \quad (41)$$

$$\bar{E}\bar{Y}\bar{X}\bar{Y} = \bar{Y}\bar{X}\bar{Y}\bar{E} = \lambda^{-1}\bar{E} \quad (42)$$

$$\bar{E}\bar{Y}^m\bar{E} = A_m\bar{E}, \text{ for all } 0 \leq m \leq k-1, \quad (43)$$

$$\sum_{l=0}^k q_l \bar{Y}^l = 0. \quad (44)$$

Furthermore Ξ is $\mathcal{B}_2^k$ -module where the actions of the generators Y, X, X^{-1} and e on Ξ are given by the maps $\bar{Y}, \bar{X}, \bar{W}$ and $\bar{E}$, respectively.

Proof. Recall that the actions of $\bar{X}, \bar{Y}$ and $\bar{E}$ on Ξ_0 are defined by the $\mathcal{B}_2^k$ -module structure on Ξ_0 . Thus all the relations hold immediately on Ξ_0 and we need only verify them on the u_{ij} and w_{ij} .

Claim: $\bar{X}\bar{W} = \bar{W}\bar{X} = 1$ and $\bar{X}\bar{E} = \bar{E}\bar{X} = \lambda\bar{E}$.

Fix integers i and j . It is easy to see that $\bar{X}$ and $\bar{E}$ preserves the three dimensional R -submodule of Ξ spanned by $\{u_{ij}, w_{ij}, \bar{E}u_{ij}\}$, and that their actions on this R -submodule are given by the matrices

$$\bar{X}' = \begin{pmatrix} 0 & 1 & 0 \\ 1 & \delta & 0 \\ 0 & -\delta\lambda & \lambda \end{pmatrix} \quad \text{and} \quad \bar{E}' = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & \lambda & A_0 \end{pmatrix},$$

respectively. Using the fact that $\lambda - \delta + \delta A_0 = \lambda^{-1}$ in R , a direct computation shows $\bar{X}'(\bar{X}' - \delta I_3 + \delta \bar{E}') = (\bar{X}' - \delta I_3 + \delta \bar{E}')\bar{X}' = I_3$. Thus $\bar{X}\bar{W}$ and $\bar{W}\bar{X}$ both act as the identity on this submodule, and in particular on u_{ij} and w_{ij} . This proves (39). Similarly, it is easily shown that $\bar{X}'\bar{E}' = \bar{E}'\bar{X}' = \lambda\bar{E}'$, so that $\bar{X}\bar{E} = \bar{E}\bar{X} = \lambda\bar{E}$ on Ξ , giving (40).

Now that we have proved (39), we may rewrite $\bar{W}$ as $\bar{X}^{-1}$. Our next step is to show that relation (41) holds on the u_{ij} .

Claim: $(\bar{X}\bar{Y}\bar{X}\bar{Y})u_{ij} = (\bar{Y}\bar{X}\bar{Y}\bar{X})u_{ij}$.

We have

$$\begin{aligned} \bar{X}\bar{Y}\bar{X}\bar{Y}u_{ij} &\stackrel{(36)}{=} \bar{X}\bar{Y}\bar{X}u_{i+1,j} \\ &\stackrel{(34)}{=} \bar{X}\bar{Y}w_{i+1,j} \\ &\stackrel{(37)}{=} \bar{X}(\bar{X}^{-1}u_{i+1,j+1} + \delta u_{1,i+j+1} - \delta YXv_{i+1,j}) \\ &\stackrel{(34),(31)}{=} u_{i+1,j+1} + \delta w_{1,i+j+1} - \delta XYXYv_{ij}. \end{aligned}$$

On the other hand,

$$\begin{aligned}
\overline{Y}\overline{X}\overline{Y}\overline{X}u_{ij} &\stackrel{(34)}{=} \overline{Y}\overline{X}\overline{Y}w_{ij} \\
&\stackrel{(37)}{=} \overline{Y}\overline{X}(\overline{X}^{-1}u_{i,j+1} + \delta u_{1,i+j} - \delta YXv_{ij}) \\
&\stackrel{(34)}{=} \overline{Y}(u_{i,j+1} + \delta w_{1,i+j} - \delta XYXv_{ij}) \\
&\stackrel{(36)}{=} u_{i+1,j+1} + \delta \overline{Y}w_{1,i+j} - \delta YXYXv_{ij} \\
&\stackrel{(38),(3)}{=} u_{i+1,j+1} + \delta w_{1,i+j+1} - \delta XYXYv_{ij}.
\end{aligned}$$

This proves our claim:

$$(\overline{X}\overline{Y}\overline{X}\overline{Y})u_{ij} = (\overline{Y}\overline{X}\overline{Y}\overline{X})u_{ij}. \quad (45)$$

To prove $(\overline{X}\overline{Y}\overline{X}\overline{Y})w_{ij} = (\overline{Y}\overline{X}\overline{Y}\overline{X})w_{ij}$, we need to first prove (42).

Claim: $\overline{Y}\overline{X}\overline{Y}\overline{E}u_{ij} = \lambda^{-1}\overline{E}u_{ij} = \overline{E}\overline{Y}\overline{X}\overline{Y}u_{ij}$ and $\overline{Y}\overline{X}\overline{Y}\overline{E}w_{ij} = \lambda^{-1}\overline{E}w_{ij} = \overline{E}\overline{Y}\overline{X}\overline{Y}w_{ij}$.

It is easy to see that

$$\begin{aligned}
\overline{Y}\overline{X}\overline{Y}\overline{E}u_{ij} &\stackrel{(32)}{=} YXYv_{00}Y^iXY^j \\
&= \lambda^{-1}v_{00}Y^iXY^j \\
&\stackrel{(32)}{=} \lambda^{-1}\overline{E}u_{ij}.
\end{aligned}$$

Recall that YXY acts as $X - \delta + \delta e$ on V . Therefore if $0 \leq i \leq k-1$, then

$$\begin{aligned}
\overline{E}\overline{Y}\overline{X}\overline{Y}u_{ij} &\stackrel{(36)}{=} \overline{E}\overline{Y}\overline{X}u_{i+1,j} \\
&\stackrel{(34)}{=} \overline{E}\overline{Y}w_{i+1,j} \\
&\stackrel{(37)}{=} \overline{E}(\overline{X}^{-1}u_{i+1,j+1} + \delta u_{1,i+j+1} - \delta YXv_{i+1,j}) \\
&\stackrel{(40)}{=} \lambda^{-1}\overline{E}u_{i+1,j+1} + \delta \overline{E}u_{1,i+j+1} - \delta eYXv_{i+1,j} \\
&\stackrel{(32)}{=} \lambda^{-1}v_{00}Y^{i+1}XY^{j+1} + \delta v_{00}YXY^{i+j+1} - \delta eYXv_{i+1,j} \\
&\stackrel{(31)}{=} \lambda^{-1}v_{00}Y^i(X - \delta + \delta e)Y^j + \delta \lambda^{-1}v_{00}Y^{i+j} - \delta eYXYv_{ij} \\
&\stackrel{(5)}{=} \lambda^{-1}v_{00}Y^iXY^j + \delta \lambda^{-1}v_{00}Y^ieY^j - \delta \lambda^{-1}ev_{ij} \\
&\stackrel{(32)}{=} \lambda^{-1}\overline{E}u_{ij} + \delta \lambda^{-1}v_{0i}eY^j - \delta \lambda^{-1}A_iv_{0j} \\
&= \lambda^{-1}\overline{E}u_{ij} + \delta \lambda^{-1}A_iv_{00}Y^j - \delta \lambda^{-1}A_iv_{00}Y^j \\
&= \lambda^{-1}\overline{E}u_{ij}.
\end{aligned}$$

Hence $\bar{Y}\bar{X}\bar{Y}\bar{E}u_{ij} = \bar{E}\bar{Y}\bar{X}\bar{Y}u_{ij} = \lambda^{-1}\bar{E}u_{ij}$. Moreover, this implies

$$\bar{Y}\bar{X}\bar{Y}\bar{E}w_{ij} \stackrel{(33)}{=} \lambda(\bar{Y}\bar{X}\bar{Y}\bar{E}u_{ij}) = \lambda\lambda^{-1}\bar{E}u_{ij} = \lambda^{-1}\bar{E}w_{ij}$$

and

$$\begin{aligned} \bar{E}\bar{Y}\bar{X}\bar{Y}w_{ij} &\stackrel{(34)}{=} \bar{E}\bar{Y}\bar{X}\bar{Y}\bar{X}u_{ij} \\ &\stackrel{(45)}{=} \bar{E}\bar{X}\bar{Y}\bar{X}\bar{Y}u_{ij} \\ &\stackrel{(40)}{=} \bar{X}\bar{E}\bar{Y}\bar{X}\bar{Y}u_{ij} \\ &= \lambda^{-1}\bar{X}\bar{E}u_{ij} \\ &\stackrel{(40)}{=} \lambda^{-1}\bar{E}\bar{X}u_{ij} \\ &\stackrel{(34)}{=} \lambda^{-1}\bar{E}w_{ij}. \end{aligned}$$

Hence $\bar{Y}\bar{X}\bar{Y}\bar{E}w_{ij} = \lambda^{-1}\bar{E}w_{ij} = \bar{E}\bar{Y}\bar{X}\bar{Y}w_{ij}$, finishing our proof of (42). We are now ready to show that $\bar{X}\bar{Y}\bar{X}\bar{Y}w_{ij} = \bar{Y}\bar{X}\bar{Y}\bar{X}w_{ij}$. By (39) and (40), we have $\bar{X}^2 = 1 + \delta\bar{X} - \delta\lambda\bar{E}$. Thus

$$\begin{aligned} \bar{Y}\bar{X}\bar{Y}\bar{X}w_{ij} &\stackrel{(34)}{=} \bar{Y}\bar{X}\bar{Y}\bar{X}^2u_{ij} \\ &= \bar{Y}\bar{X}\bar{Y}(1 + \delta\bar{X} - \delta\lambda\bar{E})u_{ij} \\ &\stackrel{(45),(42)}{=} (1 + \delta\bar{X} - \delta\lambda\bar{E})\bar{Y}\bar{X}\bar{Y}u_{ij} \\ &= \bar{X}^2\bar{Y}\bar{X}\bar{Y}u_{ij} \\ &\stackrel{(45)}{=} \bar{X}\bar{Y}\bar{X}\bar{Y}\bar{X}u_{ij} \\ &\stackrel{(34)}{=} \bar{X}\bar{Y}\bar{X}\bar{Y}w_{ij}. \end{aligned}$$

Hence we have proved (41).

For any integer l and for $0 \leq m \leq k-1$, we have

$$\bar{E}\bar{Y}^m v_{0l} = eY^m v_{0l} = ev_{ml} = A_m v_{0l}.$$

Thus $\bar{E}\bar{Y}^m$ acts as A_m on $\langle v_{0l} \mid 0 \leq l \leq k-1 \rangle$. However, it is clear from the definitions that the image of $\bar{E}$ lies in $\langle v_{0l} \mid 0 \leq l \leq k-1 \rangle$. Therefore $\bar{E}\bar{Y}^m \bar{E} = A_m \bar{E}$, giving (43).

It now remains to show that $\bar{Y}$ satisfies the k -th order relation (44). It follows directly from (30), (31) and (36) that

$$\sum_{l=0}^k q_l \bar{Y}^l v_{ij} = \sum_{l=0}^k q_l \bar{Y}^l u_{ij} = 0,$$

for all integers i and j . We shall prove $\sum_{l=0}^k q_l \bar{Y}^l w_{ij} = 0$ for all $i \geq 1$ by induction on i .

It is clear from (38) that $\bar{Y}^l w_{1j} = w_{1,j+l}$ for any integer j . Thus

$$\sum_{l=0}^k q_l \bar{Y}^l w_{1j} = \sum_{l=0}^k q_l w_{1,j+l} \stackrel{(30)}{=} 0,$$

proving the case $i = 1$. Suppose the inductive hypothesis holds for some $i \geq 1$. Recall, by definition (35), $\bar{X}w_{ij} = u_{ij} + \delta w_{ij} - \delta\lambda\bar{E}u_{ij}$. Then, using (36) and (37),

$$\bar{Y}\bar{X}w_{ij} = u_{i+1,j} + \delta\bar{X}^{-1}u_{i,j+1} + \delta^2u_{1,i+j} - \delta^2YXv_{ij} - \delta\lambda\bar{Y}\bar{E}u_{ij}.$$

Applying $\bar{X}$ to both sides and rearranging gives

$$w_{i+1,j} = \bar{X}\bar{Y}\bar{X}w_{ij} - \delta u_{i,j+1} - \delta^2w_{1,i+j} + \delta^2XYXv_{ij} + \delta\lambda\bar{X}\bar{Y}\bar{E}u_{ij}.$$

Now, the terms $u_{i,j+1}$, $XYXv_{ij}$ and $\bar{X}\bar{Y}\bar{E}u_{ij}$ lie in the R -submodule $\Xi_0 \oplus \Xi_1$. We know the operator $\sum_{l=0}^k q_l \bar{Y}^l$ acts as the zero operator on this submodule, so applying this operator to both sides gives

$$\begin{aligned} \sum_{l=0}^k q_l \bar{Y}^l w_{i+1,j} &= \sum_{l=0}^k q_l \bar{Y}^l \bar{X}\bar{Y}\bar{X}w_{ij} - \delta^2 \sum_{l=0}^k q_l \bar{Y}^l w_{1,i+j} \\ &\stackrel{(41)}{=} \bar{X}\bar{Y}\bar{X} \sum_{l=0}^k q_l \bar{Y}^l w_{ij} - \delta^2 \sum_{l=0}^k q_l \bar{Y}^l w_{1,i+j} \\ &= 0, \end{aligned}$$

by the inductive hypothesis and the $i = 1$ case proved above. Therefore we have proved by induction on i that $\sum_{l=0}^k q_l \bar{Y}^l w_{ij} = 0$, for all $i \geq 1$. However, $\{w_{ij} \mid 1 \leq i \leq k\}$ spans Ξ_2 , so (44) holds on all of Ξ . This completes the proof of the theorem. $\square$

We have now constructed a free $\mathcal{B}_2^k(R)$ -module of rank $3k^2$. The following theorem shows that Ξ is in fact isomorphic to $\mathcal{B}_2^k$, naturally considered as a $\mathcal{B}_2^k$ -module. Moreover, it states explicitly the bijection between our basis of Ξ and a basis of $\mathcal{B}_2^k$.

Theorem 4.2. (cf. Proposition 29 of Haring-Oldenburg [4])
The R -module homomorphism $\phi : \Xi \rightarrow \mathcal{B}_2^k$ defined by

$$\begin{aligned} v_{ij} &\mapsto Y^i e Y^j \\ u_{ij} &\mapsto Y^i X Y^j \\ w_{ij} &\mapsto X Y^i X Y^j \end{aligned}$$

is a $\mathcal{B}_2^k$ -module isomorphism. Furthermore $\mathcal{B}_2^k$ is R -free of rank $3k^2$ with basis $\{Y^i e Y^j, Y^i X Y^j, X Y^i X Y^j \mid i, j = 0, \dots, k-1\}$.

Proof. To show that ϕ is a $\mathcal{B}_2^k$ -module homomorphism, it suffices to check that $\phi(ax_{ij}) = a\phi(x_{ij})$ for $a \in \{X, Y, e\}$ and $x \in \{u, v, w\}$. This is

straightforward to verify using the algebra relations and the definitions of $\overline{X}$, $\overline{Y}$ and $\overline{E}$.

Let $\kappa = w_{00} - \delta u_{00} + \delta \lambda v_{00} \in \Xi$. Consider the $\mathcal{B}_2^k$ -module homomorphism $\zeta : \mathcal{B}_2^k \rightarrow \Xi$ defined by

$$a \mapsto a\kappa.$$

We will show that ϕ and ζ are mutual inverses. Certainly $\zeta(1) = \kappa$ and

$$\phi(\kappa) = X^2 - \delta X + \delta \lambda e = 1.$$

Thus $\phi\zeta(1) = 1$. But $\phi\zeta$ is a $\mathcal{B}_2^k$ -module homomorphism and 1 generates $\mathcal{B}_2^k$ as a $\mathcal{B}_2^k$ -module, so $\phi\zeta$ is the identity. The same argument will show that $\zeta\phi$ is the identity, provided we can show that κ generates Ξ , as a $\mathcal{B}_2^k$ -module. Let $\mathcal{B}_2^k\kappa$ be the submodule of Ξ generated by κ . Certainly $\mathcal{B}_2^k\kappa$ contains

$$\begin{aligned} X\kappa &\stackrel{(34),(35)}{=} (u_{00} + \delta w_{00} - \delta \lambda e u_{00}) - \delta w_{00} + \delta \lambda X v_{00} \\ &\stackrel{(32)}{=} u_{00} - \delta \lambda v_{00} X + \delta \lambda^2 v_{00} \\ &= u_{00} - \delta \lambda^2 v_{00} + \delta \lambda^2 v_{00} \\ &= u_{00}. \end{aligned}$$

Thus $\mathcal{B}_2^k\kappa$ also contains $XYu_{00} = w_{10}$. Now, for any integers i and j , the element

$$Y^{i-1}X^{-1}Y^jw_{10} \stackrel{(38)}{=} Y^{i-1}X^{-1}w_{1j} \stackrel{(34)}{=} Y^{i-1}u_{1j} \stackrel{(36)}{=} u_{ij}$$

is contained in $\mathcal{B}_2^k\kappa$. Applying X , we also have $w_{ij} \in \mathcal{B}_2^k\kappa$. Finally $\mathcal{B}_2^k\kappa$ contains

$$\begin{aligned} \lambda^{-1}Y^ie u_{0j} &\stackrel{(32)}{=} \lambda^{-1}Y^iv_{00}XY^j \\ &= Y^iv_{00}Y^j \\ &= v_{ij} \end{aligned}$$

for all integers i and j . Therefore $\mathcal{B}_2^k\kappa = \Xi$, so that ϕ and ζ are mutual inverses as claimed. Hence $\mathcal{B}_2^k(R)$ is a free R -module of rank $3k^2$ and $\{Y^ieY^j, Y^iXY^j, XY^iXY^j \mid i, j = 0, \dots, k-1\}$ is an R -basis of $\mathcal{B}_2^k(R)$. $\square$

Corollary 1. (Lemma 26 of Häring-Oldenburg [4])

The two-sided ideal of $\mathcal{B}_2^k(R)$ generated by e is a free R -module with basis $\{Y^ieY^j \mid i, j = 0, \dots, k-1\}$. Moreover, as a $\mathcal{B}_2^k(R)$ -module, it is isomorphic to a direct sum decomposition of k copies of V .

Proof. By Theorem 4.2, the restriction of ϕ to $\Xi_0 = V \otimes_R V$ is an injective $\mathcal{B}_2^k$ -module homomorphism $\phi' : V \otimes_R V \hookrightarrow \mathcal{B}_2^k$. Now $V \otimes_R V$ has an R -basis $\{v_{ij} \mid i, j = 0, \dots, k-1\}$, so

$$\{\phi'(v_{ij}) \mid i, j = 0, \dots, k-1\} = \{Y^ieY^j \mid i, j = 0, \dots, k-1\}$$

is an R -basis for $\text{Im}(\phi')$. Now $\text{Im}(\phi')$ is a left ideal of $\mathcal{B}_2^k$, since ϕ' is a $\mathcal{B}_2^k$ -module homomorphism. But $(Y^i e Y^j)^* = Y^j e Y^i$. Thus $\text{Im}(\phi')$ is invariant under $*$, so it is a two sided ideal of $\mathcal{B}_2^k$. It clearly contains and is generated by e , so it is exactly the two sided ideal generated by e . Finally under ϕ' , it is isomorphic as a $\mathcal{B}_2^k$ -module to

$$V \otimes_R V = \bigoplus_{i=0}^{k-1} V \otimes_R v_i,$$

that is, a direct sum of k copies of V . □

REFERENCES

- [1] S. Ariki and K. Koike. A Hecke algebra of $(\mathbb{Z}/r\mathbb{Z}) \wr \mathfrak{S}_n$ and construction of its irreducible representations. *Adv. Math.*, 106:216–243, 1994.
- [2] J. S. Birman and H. Wenzl. Braids, link polynomials and a new algebra. *Trans. Amer. Math. Soc.*, 313:249–273, 1989.
- [3] F. Goodman and H. Hauschild. Affine Birman-Wenzl-Murakami algebras and tangles in the solid torus. *Fund. Math.*, 190:77–137, 2006.
- [4] R. Häring-Oldenburg. Cyclotomic Birman-Murakami-Wenzl algebras. *J. Pure Appl. Algebra*, 161:113–144, 2001.
- [5] J. Murakami. The Kauffman polynomial of links and representation theory. *Osaka J. Math.*, 24:745–758, 1987.

STEWART WILCOX, DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY,
1 OXFORD STREET, CAMBRIDGE MA 02318, USA
E-mail address: `swilcox@fas.harvard.edu`

SHONA YU, SCHOOL OF MATHEMATICS AND STATISTICS F07, UNIVERSITY OF
SYDNEY, SYDNEY NSW 2006, AUSTRALIA
E-mail address: `shonayu@maths.usyd.edu.au`